

Measurement of direct neutron capture by neutron-rich sulfur isotopes

H. Beer^a, C. Coceva^b, R. Hofinger^c, P. Mohr^c, H. Oberhummer^c, P.V. Sedyshev^d and Yu.P. Popov^d

^aForschungszentrum Karlsruhe, IK III, P.O. Box 3640, D-76021 Karlsruhe, Germany

^bENEA, Via Don Fiammelli 2, I-40128 Bologna, Italy

^cInstitut für Kernphysik, TU Wien, Wiedner Hauptstraße 8–10, A-1040 Wien, Austria

^dFrank Laboratory, JINR, 141980–Dubna, Moscow Region, Russia

Thermal neutron capture cross sections for $^{34}\text{S}(n,\gamma)^{35}\text{S}$ and $^{36}\text{S}(n,\gamma)^{37}\text{S}$ have been measured and spectroscopic factors of the final states have been extracted. The calculated direct-capture cross sections reproduce the experimental data.

1. INTRODUCTION

Reaction rates of the neutron-rich S-isotopes are of interest in the nucleosynthesis of nuclei in the s-process in the S–Cl–Ar–Ca region, inhomogeneous big-bang scenario, and in the α -rich freeze out of the neutron-rich hot neutrino bubble in supernovae of type II. In the last years the importance of the direct capture (DC) has been realized. DC dominates over the compound-nucleus reaction mechanism if there exist no compound-nucleus levels near the threshold that can be excited in the reaction. This is often the case for neutron capture for neutron-rich target nuclei at the border of stability. With respect to astrophysical applications thermal measurements of such capture reactions are an important supplement to measurements in the thermonuclear energy region. Such measurements allow a more reliable extrapolation to lower energies. Furthermore, when there are no resonant contributions at thermal energy, experimental thermal cross sections allow the extraction of spectroscopic factors. These spectroscopic factors are needed to calculate the DC contribution at thermonuclear energies. This extraction is not as much model-dependent as the extraction of spectroscopic factors from direct transfer reactions, like (d,p), (^3He , ^4He), or other transfer reactions at higher energies (about 15–40 MeV). This is mainly due to the fact that the neutron optical potentials necessary for the extraction of the spectroscopic information from experimental thermal (n, γ)-data is better determined than the optical proton and especially light-ion potentials at higher energies (because for higher energies there is a larger absorption).

The purpose of this work is to show that for thermal neutron capture by target isotopes close to the border of stability, like ^{34}S and ^{36}S , the DC mechanism dominates and measurements of capture cross sections at thermal energies can be used to extract spectroscopic factors for the calculation of DC cross sections at thermonuclear energies.

Table 1

The intensity per incident neutron, I_γ , and statistical uncertainty, ΔI_γ , of the γ -ray transitions with initial and final spin/parity, J_i^Π and J_f^Π , respectively, and the transition energy E_γ . The partial and total capture cross sections of $^{34,36}\text{S}$, σ^p and σ , were determined.

Target Isotope	E_γ^a (keV)	Transition $J_i^\Pi \rightarrow J_f^\Pi$	$I_\gamma \pm \Delta I_\gamma$ 10^{-5}	σ^p (mbarn)	σ (mbarn)
^{34}S	2022.9 ^p	$1/2^+ \rightarrow 3/2^-$	50.73 ± 0.91	26.9 ± 1.1	
	2082.6 ^p	$1/2^+ \rightarrow 1/2^-$	70.34 ± 0.98	37.2 ± 1.4	
	2796.8 ^p	$1/2^+ \rightarrow 1/2^-$	27.65 ± 0.77	14.6 ± 0.7	
	3184.0 ^p	$1/2^+ \rightarrow 3/2^-$	31.99 ± 0.93	16.9 ± 0.8	
	4638.2 ^p	$1/2^+ \rightarrow 3/2^-$	280.82 ± 2.53	148.6 ± 5.5	244.3 ± 8.8
	1572.4 ^s	$1/2^+ \rightarrow 3/2^+$	179.76 ± 1.43		
	2347.8 ^s	$3/2^- \rightarrow 3/2^+$	225.76 ± 158		
	3802.0 ^s	$3/2^- \rightarrow 3/2^+$	14.43 ± 0.79		
	4189.3 ^s	$1/2^- \rightarrow 3/2^+$	15.37 ± 0.88		
	4903.4 ^s	$1/2^- \rightarrow 3/2^+$	23.49 ± 1.03		
	4963.1 ^s	$3/2^- \rightarrow 3/2^+$	12.19 ± 0.80		249.3 ± 9.0
				average	247 ± 9
^{36}S	1665.7 ^p	$1/2^+ \rightarrow 1/2^-$	6.67 ± 0.77	54.8 ± 6.9	
	2311.6 ^p	$1/2^+ \rightarrow 3/2^-$	3.81 ± 0.59	31.3 ± 5.1	
	3657.3 ^p	$1/2^+ \rightarrow 3/2^-$	18.66 ± 0.80	153.3 ± 9.9	239 ± 15
	646.2 ^s	$3/2^- \rightarrow 7/2^-$	30.37 ± 0.97		249 ± 14
	3103.3	decay line	28.77 ± 0.95		236 ± 14
				average	242 ± 12

^a Ref. [8], ^p primary transition, ^s secondary ground state transition

2. EXPERIMENT

The measurements were carried out at the reactor BR1 of the "Studiecentrum voor kernenergie" in Mol, Belgium. The setup was the same as in a previous experiment [1]. The sample of elemental sulfur (total weight 313.6 mg) contained in a teflon cylinder of 6 mm inner diameter was enriched in ^{34}S by 93.26 % and in ^{36}S by 5.933 %. Fig. 1 shows part of the γ -ray spectrum accumulated with a Ge-detector. Neutron flux and efficiency of the detector were determined in a calibration run on ^{35}Cl . The total capture cross sections of $^{34,36}\text{S}$ were calculated from the sum of primary and secondary ground state transitions and the ^{37}S γ -decay line (Table 1). The data were also corrected for multiple neutron scattering (^{34}S : 5.6 %, ^{36}S : 7 %). For the $^{34,36}\text{S}$ cross sections systematic uncertainties of 3.6 and 4.8 %, respectively, were estimated. Our thermal ^{34}S capture cross section is in good agreement with the recommended value [2]. Our ^{36}S result is consistent with the previous measurement of Raman et al. [3].

3. DIRECT-CAPTURE CALCULATIONS

The DC formalism including the folding procedure used in this work is described in Ref. [4]. The DC calculations are performed in the same manner as given already for




Figure 1. Intensities of ^{35}S and ^{37}S γ -lines

$^{36}\text{S}(n,\gamma)^{37}\text{S}$ in [5]. The input data for the folding potentials are: experimental charge distributions [6], thermal elastic scattering data [7], excitation energies of the final states [8]. For the DC-calculations the input data for the masses and Q -values are taken from Ref. [9]. The spectroscopic factors are listed in Table 2 and are taken from (d,p) -reactions [8] (except the (d,p) -data from [10] which has been taken from the original work, see Note added in Proof in [5]). The resulting thermal cross sections calculated in the DC-model are compared with the experimental cross sections for $^{34}\text{S}(n,\gamma)^{35}\text{S}$ and $^{36}\text{S}(n,\gamma)^{37}\text{S}$ in the last two columns of Table 2.

4. DISCUSSION

The DC-mechanism reproduces not only the summed thermal experimental cross sections of $^{34}\text{S}(n,\gamma)^{35}\text{S}$ and $^{36}\text{S}(n,\gamma)^{37}\text{S}$, but also the transitions to the different relevant final states. This is certainly not true for statistical model calculations [11–13], because the statistical approach is valid only in the case of a high-level density in the populated excitation range of compound nuclei. Hauser-Feshbach calculations for the above reactions lead to a significant overestimation of the cross sections by about one order of magnitude [14].

The spectroscopic factors extracted from (d,p) -reactions by different groups lead to calculated thermal cross sections that are uncertain by at least $\pm 20\%$ (see last but one column in Table 2). Spectroscopic factors extracted from thermal experimental (n,γ) -reactions can be determined more accurately, because the optical potentials are much better known in this case. Therefore, thermal (n,γ) -measurements are an excellent alternative to (p,d) - or other transfer reaction measurements in determining spectroscopic factors. These are necessary for calculating the non-resonant direct part of the cross section at the astrophysically relevant thermonuclear energies.

Table 2

Spin/parity assignments J^Π , excitation energies E_x , and neutron spectroscopic factors of states in ^{35}S and ^{37}S . The spectroscopic factors $S(\text{d,p})$ and $S(\text{n},\gamma)$ are extracted from experimental (d,p)-reactions and from the thermal neutron cross sections measured in this work, respectively. The calculated σ^{cal} and experimental σ^{exp} thermal capture cross sections are compared for $^{34}\text{S}(\text{n},\gamma)^{35}\text{S}$ and $^{36}\text{S}(\text{n},\gamma)^{37}\text{S}$. The sum of the contributions for all other transitions not shown in this table give less than $1\,\mu\text{barn}$.

J^Π	E_x (MeV)	$S(\text{d,p})$	$S(\text{n},\gamma)$	σ^{cal} (mbarn)	σ^{exp} (mbarn)
$^{34}\text{S}(\text{n},\gamma)^{35}\text{S}$					
$3/2^-$	2.348	0.32–0.55	0.48	99–169	149
$3/2^-$	3.802	0.09	0.08	20	17
$1/2^-$	4.189	0.12–0.14	0.15	12–14	15
$1/2^-$	4.903	0.44–0.80	0.49	33–60	37
$3/2^-$	4.963	0.18–0.22	0.19	26–32	27
sum				190–295	245
$^{36}\text{S}(\text{n},\gamma)^{37}\text{S}$					
$3/2^-$	0.647	0.44–0.70	0.54	126–200	153
$3/2^-$	1.992	0.04–0.08	0.19	7–13	31
$1/2^-$	2.638	0.48–0.80	0.92	29–48	55
sum				162–261	239

REFERENCES

1. H. Beer, C. Coceva, P.V. Sedyshev, Yu.P. Popov, H. Herndl, R. Hofinger, P. Mohr, and H. Oberhummer, Phys. Rev. C, submitted.
2. E. Gryntakis, D.E. Cullen and G. Mundy, IAEA, Vienna, report 273 (1987) 199.
3. S. Raman, R.F. Carlton, J.C. Wells, E.T. Jurney and J.E. Lynn, Phys. Rev. C 32 (1985) 18.
4. E. Krausmann, W. Balogh, H. Oberhummer, T. Rauscher, W. Ziegert and K.-L. Kratz, Phys. Rev. C 53 (1996) 469.
5. H. Beer, P.V. Sedyshev, Yu.P. Popov, W. Balogh, H. Herndl and H. Oberhummer, Phys. Rev. C 52 (1995) 3442.
6. H. De Vries, C.W. De Jager, and C. De Vries, At. Data and Nucl. Tables 36 (1987) 495.
7. V.F. Sears, Neutron News 3 (1992) 26.
8. P.M. Endt, Nucl. Phys. A 521 (1990) 1.
9. G. Audi and A.H. Wapstra, Nucl. Phys. A 565 (1993) 1.
10. G. Eckle, H. Kader, H. Clement, F. J. Eckle, F. Merz, R. Hertenberger, H. J. Maier, P. Schiemenz and G. Graw, Nucl. Phys. A 491 (1989) 205.
11. R.K. Wallace, S.E. Woosely and T.A. Weaver, ApJ 258 (1981) 696.
12. F.K. Thielemann, M. Arnould and J.J. Truran, Advances in Nuclear Astrophysics, Editions Frontières, Gif-sur-Yvette, 1987, p. 525.
13. J.J. Cowan, F.K. Thielemann and J.W. Truran, Phys. Rep. 208 (1991) 268.
14. T. Rauscher, private communication (1996).